

Mathematics for Materials

Problem Set A for 'Vectors, Matrices...' Course

(overall 3rd set in the Michaelmas Term maths provision)

These first 4 questions should be do-able after the first lecture of the course.

1. For each of the following pairs of points A and B find:
 - (i) The vector from A to B , $\overrightarrow{AB}$ in $\begin{pmatrix} x \\ y \\ z \end{pmatrix}$ form
 - (ii) The vector $\overrightarrow{AB}$ in $\hat{i}, \hat{j}, \hat{k}$ form
 - (iii) The magnitude, $|\overrightarrow{AB}|$
 - (iv) The unit vector of $\overrightarrow{AB}$
 - (v) The direction cosines of $\overrightarrow{AB}$

(a) $A = (0, 0, 0)$ and $B = (2, 3, 0)$. (b) $A = (1, -3, 2)$ and $B = (0, 1, 0)$.
(c) $A = (-1, -1, 3)$ and $B = (2, 4, -2)$.
2. Consider the regular hexagon $OABCDE$ with O at the origin, where A has position vector $\mathbf{a}$, and B has position vector $\mathbf{a} + \mathbf{b}$.
 - (a) Express the position of the other three vertices in terms of $\mathbf{a}$ and $\mathbf{b}$. *Hint: Draw a sketch and solve the problem graphically.*
 - (b) If someone moves around the hexagon (from O to A to B etc.), what are the expressions for each of the six moves they make in terms of $\mathbf{a}$ and $\mathbf{b}$?
3. You are out walking when one of the other people in the group is injured and needs medical assistance. Consulting a map, you decide that you are at 5 km east, 3 km south (relative to the centre of the map), and at an elevation of 200 m. The two closest towns are Rushton at 1 km east, 3 km north, elevation 800 m and Charmoor at 2 km west, 1 km south, elevation 500 m. Clearly showing your reasoning, answer the following:
 - (a) What compass bearing should you follow to reach each town?
 - (b) Assuming you walk in a straight line, which town is closer?
 - (c) If one kilometre of net vertical change costs you five times as much energy as one kilometre of lateral movement, which town can you reach more efficiently?
4. In triangle ABC , D is the mid-point of the side AB , E is the mid-point of side BC , and X is the intersection of CD and AE . Using vectors, find the ratio of the length of CX to the length of XD and similarly find the ratio of AX to XE . *Hints: Define two vectors $\mathbf{u}$ and $\mathbf{v}$ to coincide with two sides of the triangle, and then express everything else in terms of those vectors. The ratio you are looking for is a pure number, not depending on the lengths of the sides (so it's the same for all triangles) and point X divides CD in the same ratio as it divides AE . Lookup "centroid" for more information.*

The remaining questions should be do-able after the first week of the course is complete

5. (a) Use a diagram to explain why $\mathbf{r} = \lambda \mathbf{a} + \mathbf{b}$ describes a line.
 (b) Find the vector equation of the line containing the points $(2, -2, 5)$ and $(0, -3, 1)$.
 (c) Find the vector equation of the line containing the points $2\mathbf{u} - \mathbf{v} + 3\mathbf{w}$ and $-3\mathbf{u} + \mathbf{w}$ where $\mathbf{u}$, $\mathbf{v}$, and $\mathbf{w}$ are non-coplanar vectors. *N.B. You should not assume that $\mathbf{u}$, $\mathbf{v}$, and $\mathbf{w}$ are orthogonal or that they are unit vectors.*
6. (a) Use a diagram to explain why $\mathbf{r} = \lambda \mathbf{a} + \mu \mathbf{b} + \mathbf{c}$ describes a plane.
 (b) Find the vector equation of the plane containing the points $(3, -1, 3)$, $(1, 1, -2)$, and $(0, 2, 4)$.
 (c) A plane contains two points whose position vectors from the origin are $\overrightarrow{OA} = \mathbf{u} + \mathbf{v} + \mathbf{w}$ and $\overrightarrow{OB} = 2\mathbf{u} - \mathbf{v} - \mathbf{w}$. Furthermore, the vector $-\mathbf{u} + 2\mathbf{v}$ is parallel to the plane. Find the vector equation for this plane, given that $\mathbf{u}$, $\mathbf{v}$, and $\mathbf{w}$ are non-coplanar vectors.
7. (a) Using a sketch, explain the geometrical meaning of the equation of the dot product, $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos \theta$.
 (b) Explain how you can use the dot product to find the part of $\mathbf{a}$ parallel to $\mathbf{b}$.
 (c) Resolve the vector $\begin{pmatrix} 1 \\ -3 \\ 4 \end{pmatrix}$ into two vectors, one parallel to $\begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$ and one perpendicular.
8. (a) Use a diagram to explain why $\mathbf{r} \cdot \mathbf{n} = d$ describes a plane. What meaning does d have if $\mathbf{n}$ is a unit vector?
 (b) A plane contains the point $(3, -1, 3)$, and vector $\mathbf{n} = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$ is normal to the plane. What is the shortest distance to point $(4, 3, 1)$?
9. (a) Write down and explain the vector equation of a circle using a diagram. Explain how this relates to the Cartesian equation for a circle.
 (b) Write down the vector and Cartesian equations of a sphere, centre $(1, 2, -4)$ and radius 5.
10. (a) Vectors $\mathbf{a}$ and $\mathbf{b}$ are orthogonal. What are $\mathbf{a} \cdot \mathbf{b}$, $\mathbf{a} \cdot \mathbf{a}$, $|\mathbf{a} \times \mathbf{b}|$, and $|\mathbf{b} \times \mathbf{b}|$?
 (b) Let the components of $\mathbf{a}$ and $\mathbf{b}$ be a_x, a_y, a_z and b_x, b_y, b_z respectively. Show that $\mathbf{a} \times \mathbf{b}$ is perpendicular to $\mathbf{a}$ and to $\mathbf{b}$.
11. (a) Using a sketch, explain the geometrical meaning of the equation of the cross product, $|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}||\mathbf{b}| \sin \theta$.
 (b) Using vectors, find the area of a parallelogram having one corner at the origin $(0, 0, 0)$ and the adjacent two corners at $(1, 0, 3)$ and $(4, -2, -2)$.
 (c) Using vectors, find the area of the triangle with corners at $(0, 0, 0)$, $(-2, 3, 2)$ and $(1, 1, -4)$.
12. (a) Transform the plane equation $\mathbf{r} = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 0 \\ -1 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$ into the form $\mathbf{r} \cdot \mathbf{n} = d$, and hence find the distance to the point $(3, 5, 7)$. Explain your working.
 (b) Do the lines $\mathbf{r} = \begin{pmatrix} 5 \\ 1 \\ 2 \end{pmatrix} + \lambda \begin{pmatrix} 4 \\ 1 \\ 1 \end{pmatrix}$ and $\mathbf{r} = \begin{pmatrix} -3 \\ 1 \\ -2 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 3 \\ -4 \end{pmatrix}$ define a plane? Justify your answer.
 (c) $\mathbf{a}$, $\mathbf{b}$, and $\mathbf{c}$ are non-coplanar vectors. Does the line $\mathbf{r} = 5\mathbf{a} - 8\mathbf{b} + 6\mathbf{c} + \gamma(-2\mathbf{a} - 16\mathbf{b} + 13\mathbf{c})$ lie in the plane $\mathbf{r} = 3\mathbf{a} + \mathbf{c} + \lambda(\mathbf{a} - 2\mathbf{b} + \mathbf{c}) + \mu(-\mathbf{a} - 2\mathbf{b} + 2\mathbf{c})$? Justify your answer.
13. What is the shortest distance between point $(2, 0, -3)$ and the line $\mathbf{r} = \begin{pmatrix} 4 \\ 1 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 1 \\ -2 \end{pmatrix}$?
 Use a sketch to explain your working.