

Mathematics for Materials

Problem Set D for the 'Vectors, Matrices...' Course

(overall 6th set in the Michaelmas Term maths provision)

Online Problems

The following questions 1 to 5 can be answered using the **online system**. They are included on this sheet for reference only.

1. What is the value of the determinant $\begin{vmatrix} 1 & 5 \\ -3 & 2 \end{vmatrix}$? (a) 11 (b) 17 (c) 3 (d) -13.
2. What is the value of the determinant $\begin{vmatrix} 2 & -7 \\ -7 & 2 \end{vmatrix}$? (a) 45 (b) -5 (c) 53 (d) -45.
3. What is the value of the determinant $\begin{vmatrix} 0 & 2 & 0 \\ 3 & -1 & 4 \\ 2 & 5 & 1 \end{vmatrix}$? (a) -22 (b) 22 (c) 10 (d) 0.
4. What is the value of the determinant $\begin{vmatrix} 6 & 2 & 1 \\ -1 & -1 & 1 \\ 1 & 4 & -3 \end{vmatrix}$? (a) 7 (b) 43 (c) -5 (d) -13.
5. What is the value of the determinant $\begin{vmatrix} 1 & 0 & 0 & 2 \\ -1 & 3 & 0 & 1 \\ 0 & -2 & 1 & 1 \\ 0 & 5 & -3 & 2 \end{vmatrix}$? (a) -36 (b) 18 (c) 7 (d) 14.

Hint: Remember you can add/subtract rows from one another, then evaluate the determinant.

Pen and Paper Problems

1. (a) Show that $\begin{vmatrix} a & b & c \\ d & f & g \\ h & j & k \end{vmatrix}$ represents the volume of a three dimensional shape. *N.B. Figure out the volume by some means other than a determinant, and compare!*
(b) Show that $|\mathbf{A}^T| = |\mathbf{A}|$, i.e. that $\begin{pmatrix} a & d & h \\ b & f & j \\ c & g & k \end{pmatrix}$ spans the same volume as $\begin{pmatrix} a & b & c \\ d & f & g \\ h & j & k \end{pmatrix}$.
2. Using your results from question 1, show geometrically how the determinant of a 3×3 matrix is affected by the following:
 - (a) Any two rows are swapped
 - (b) A multiple of one row is added to another row
 - (c) A row is multiplied by a constant
 - (d) A row contains only zeros
 - (e) One row is proportional to another row
 - (f) What happens to the above properties if we consider columns instead of rows?
3. Using the properties of determinants, simplify and evaluate these determinants. Do not use a calculator.

(a) $\begin{vmatrix} 30 & -150 & -45 \\ 51 & 85 & 306 \\ 77 & 0 & 462 \end{vmatrix}$

(b) $\begin{vmatrix} 4 & x & -1 \\ 8x & 4x^2 & -12xy \\ y^2 & x & 3 \end{vmatrix}$

4. (a) Evaluate

$$\begin{vmatrix} 4 & 0 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 & 0 \\ 7 & -5 & -3 & 0 & 0 \\ 16 & -4 & 8 & 2 & 0 \\ 1 & 1 & 2 & 3 & 5 \end{vmatrix}$$

- (b) The above matrix is called a *lower triangular matrix*. Its determinant can be found by what simple calculation?
- (c) Use your answer to (b) to evaluate the following (an *upper triangular matrix*):

$$\begin{vmatrix} 2 & 37 & 51 & 412 & -73 \\ 0 & 3 & 82 & -115 & 345 \\ 0 & 0 & 5 & 666 & 382 \\ 0 & 0 & 0 & 7 & 525 \\ 0 & 0 & 0 & 0 & 11 \end{vmatrix}$$

- (d) Use the properties of determinants to convert the following to an upper triangular matrix and thus evaluate its determinant:

$$\begin{vmatrix} 1 & 2 & 0 & 0 & 0 \\ 2 & 3 & 4 & 0 & 0 \\ 0 & 4 & 5 & 6 & 0 \\ 0 & 0 & 6 & 7 & 8 \\ 0 & 0 & 0 & 8 & 9 \end{vmatrix}$$

5. Find the inverse of each of the following, using either the method of cofactors or Gaussian elimination:

(a) $\begin{pmatrix} 3 & -7 \\ 5 & 1 \end{pmatrix}$

(b) $\begin{pmatrix} 2 & 3 & -2 \\ 8 & 4 & 1 \\ 6 & -3 & -5 \end{pmatrix}$

(c) $\begin{pmatrix} 3 & 1 & -1 \\ 0 & 6 & 2 \\ -1 & 4 & 9 \end{pmatrix}$

6. If the elements of a transformation matrix are constants, it transforms a cube into a parallelepiped. For a cube of volume V initially aligned along the x, y, z axis, find the volume of the parallelepiped formed after the following transformations:

(a) $\begin{pmatrix} 1 + \varepsilon_x & 0 & 0 \\ 0 & 1 + \varepsilon_y & 0 \\ 0 & 0 & 1 + \varepsilon_z \end{pmatrix}$

(b) $\begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ 0 & 0 & 1 \end{pmatrix}$

- (c) A pure shear of magnitude τ in the $x - y$ plane in the x and y directions

(d) $\begin{pmatrix} 1 & 2 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & \frac{1}{2} \end{pmatrix}$

- (e) Which types of transformation change the cube's volume, and which do not?

7. Find the matrix for the transformation that 'undoes' each of the following and explain your answers:

- (a) A rotation of θ clockwise about the y axis

- (b) Pure shear of magnitude τ on the $x - z$ plane accompanied by 50% compression in the y direction

8. Solve the following system of simultaneous equations using each of the following methods:

$$\begin{aligned}2x - y + 2z &= 6 \\4x + 3y - 7z &= -11 \\x + y - 3z &= -6\end{aligned}$$

- (a) Cramer's rule
- (b) Gaussian elimination
- (c) Finding the inverse matrix using cofactors